

Smoking and reproduction

Introduction

This fact sheet explains how smoking affects fertility in both men and women. It reviews the harm caused by smoking during pregnancy as well as some of the longer-term risks to the health of children born to parents who smoke.

A published review of the literature examining the relationship between smoking and fertility concluded that tobacco use affects every system involved in the reproductive process.¹ This finding is supported by a systematic review of the scientific literature on the impact of cigarette smoking and smoke constituents on the different stages of reproductive function which found that “all stages of reproductive functions are targets of cigarette smoke toxicants.”²

A study published by the US Centers for Disease Control and Prevention notes that while smoking rates amongst pregnant women in the Western world have fallen in recent years, smoking remains a major cause of new-born deaths, early births and babies born with low birth weight. Researchers found that smoking was associated with:

- 5-8% of premature births
- 13-19% of cases of low birth weight in babies carried to full term
- 5-7% of preterm-related deaths
- 23-34% of deaths caused by sudden infant death syndrome (cot death).³

In the UK, smoking in pregnancy causes up to 5,000 miscarriages, 300 peri-natal deaths and around 2,200 premature births each year.⁴

Female fertility

Research has established beyond doubt that smoking can have a negative impact on female fertility. Women who smoke take longer to conceive than women who do not smoke.⁵

One study found that tobacco consumption affects uterine receptivity, with heavy smokers more likely to be affected.⁶ However, even comparatively low levels of smoking can have a significant impact on female fertility. There is also a higher rate of ectopic pregnancies in smokers.^{7,8}

There is evidence to suggest that smoking reduces the success rates of fertility treatment.⁹ Studies of women undergoing assisted reproductive treatment have demonstrated a significant negative effect associated with smoking.^{10,11,12} One study in the 1990s showed a 50% reduction in implantation rate among smokers compared to women who had never smoked.¹³

A growing body of research suggests that maternal smoking may have a negative impact on the fertility of both female and male off-spring.^{14,15,16} Smoking during pregnancy reduces the number of germ cells (the cells that form eggs in females and sperm in males) and somatic cells (the cells that form every other part of the body) that form in the developing foetus.¹⁷ Smoking during

pregnancy also has an impact on protamine, a protein essential in sperm production which can lead to fertility problems.¹⁸ However, further research into this issue is needed before a causal link can be definitively established.

Male fertility & sexual impotence

Cigarette smoking affects male fertility, even starting in utero. In one animal study female mice were exposed to cigarette smoke before and after becoming pregnant. The male offspring were observed to have depleted sperm counts, damage to germ cell DNA and deviant testicular development.¹⁹ Other studies further show that men whose mothers smoked while pregnant are at risk of having smaller testes,²⁰ lower mean sperm concentration and lower total sperm counts.²¹

Men who smoke have a lower sperm count than non-smokers, and their semen contains a higher proportion of malformed sperm.^{22,23,24} By-products of nicotine present in the semen of smokers have been found to reduce the motility of sperm and their fertilization capacity.^{24,25} While there is a correlation between the number of cigarettes smoked and the damage to sperm, researchers have found that there is no “safe” level of smoking. Even light smoking is associated with reduced male fertility.^{22,26}

Mounting evidence shows a significant association between smoking and male sexual impotence with the association increasing with the number of cigarettes smoked per day.^{27,28,29,30,31}

Smoking in pregnancy

Younger mothers, women in disadvantaged circumstances and who have never worked tend to be more likely to smoke throughout their pregnancy.³²

In March 2011, as part of the Tobacco Control Plan for England, the Government set an ambition to reduce the percentage of women who smoked during pregnancy from 14% to 11% by 2015 (measured at the time of birth).³³

In response to a challenge from the then Public Health Minister to produce recommendations on how the ambition could be achieved, the Smoking in Pregnancy Challenge Group was established. This is a partnership of experts from academia, medical colleges and the third sector.³⁴

By April 2016, 10.6% of women were smoking at the time of delivery, making this the first year that annual figures have been below the 11% target. However, this still equates to just over 67,000 infants born to smoking mothers in England.³⁵

Following the attainment of the Government’s 11% target, the Smoking in Pregnancy Challenge Group has proposed a new target to reduce the percentage of women smoking during pregnancy to 6% or less by 2020.³⁶

Foetal growth and birth weight

Maternal smoking is a major risk factor for low birth weight and babies who are small for their gestational age (SGA).^{37,38,39,40} Research in Sweden has shown that babies born to women who smoke throughout their pregnancy are on average 162-226 grams lighter than babies born to non-smoking mothers.⁴¹ Spanish researchers found the mean difference in their study subjects to be 216 grams.⁴² One study reported that approximately 30% of growth-restricted neonates could be independently associated with maternal smoking.⁴³ Others found that smoking during pregnancy can lead to slower growth of the foetus’ head⁴⁴ and smaller head circumference at birth.^{45,46}

Perinatal mortality and Sudden Infant Death Syndrome

Perinatal mortality includes still-birth (loss of the foetus after the 24th week of pregnancy) and neonatal death (death of the newborn within the first four weeks of life). It is estimated that about one-third of all perinatal deaths in the UK are caused by maternal smoking.⁴⁷ This equates to approximately 300 deaths per year. Evidence from the West Midlands has demonstrated that “babies born to women who smoke during pregnancy are around 40% more likely to die within the first four weeks of life than babies born to non-smokers.”^{40,48}

More than one-quarter of the risk of death due to Sudden Infant Death Syndrome (cot death) is attributable to smoking during pregnancy and exposure to secondhand smoke, particularly in the home.^{49,50,51,52} The risk of cot death is trebled in infants whose mothers smoke both during and after pregnancy.⁵³ The greater the number of cigarettes smoked, the higher the risk of cot death.^{53,54} One possible explanation for this is that nicotine and its derivatives found in tobacco smoke easily reach the cerebrospinal fluid in the foetus, causing damage to the ependymal (the lining providing a protective barrier and filtration system separating the brain from cerebrospinal fluid).⁵⁵ One way to help reduce the risk of cot death is to maintain a smoke-free home and vehicle at all times.⁵⁶

Pre-term birth is a major clinical problem, accounting for about half of all neonatal deaths. Recent research in Sweden examined the relationship between maternal smoking and pre-term birth and found that, compared to non-smokers, moderate smokers had a two-fold increase in risk of preterm labour, which rose to two and a half times greater risk among heavy smokers.⁵⁷ Maternal smoking negatively impacts offspring from infancy to childhood and even into adulthood.

Maternal smoking and child health

Infants and children of parents who smoke are twice as likely to suffer from a serious **respiratory infection** as the children of non-smokers. Smoking during pregnancy can also increase the risk of **asthma and wheezing** in young children.^{58,59,60} The effects of maternal smoking during pregnancy and childhood environmental tobacco smoke (ETS) exposure on asthma and wheezing were investigated in 5,762 school-aged children in California. Maternal smoking during pregnancy was consistently associated with varying forms of both asthma and wheezing. In contrast current and previous ETS exposure was not associated with asthma but was associated with subcategories of wheezing.⁶¹ Research suggests that the increased risk of asthma and respiratory infections may be due to changes in biological receptors in the baby's immune system that are responsible for recognising and fighting infections and bacteria.⁶² Smoking in pregnancy is also associated with an increased risk of **infantile colic**.^{63,64}

Smoking in pregnancy may also have implications for the **long term physical growth and intellectual development** of the child.^{65,66}

There is also evidence that smoking interferes with a woman's **hormonal balance** during pregnancy and that this may have long-term consequences on the reproductive organs of her children.⁶⁷ (See fertility sections above.)

A 27 year study examining cholesterol in children found evidence suggesting that maternal smoking in pregnancy is associated with an increased rise in total **cholesterol** levels and a tendency towards an adverse lipoprotein profile in the offspring.⁶⁸ Another study concluded that smoking among pregnant women is independently associated with significantly lower high-density lipoprotein cholesterol in healthy 8-year-olds.⁶⁹ Adults who were small for gestational age at birth as a result of maternal smoking also have an increased risk of hypercholesterolemia (high cholesterol).⁷⁰

There is also evidence of a link between maternal smoking during pregnancy and risk of childhood cancer. In a case-control study researchers observed a dose-response relationship between the number of cigarettes smoked per day by the mother during pregnancy and cancer risk in offspring. Compared to a control group the risk for the experimental group was doubled for non-Hodgkin lymphoma, acute lymphoblastic leukaemia, and Wilms' tumour.⁷¹ More recent research has supported this finding that there is a link between maternal smoking and risk of childhood cancer.^{72,73}

Although confounding bias (i.e. a situation in which the effect or association between an exposure and outcome is distorted by the presence of another variable) may contribute to these risk factors,⁷⁴ further impacts include:

- Increased risk of **congenital defects** in the offspring of smokers,⁷⁵ including orofacial clefts;^{76,77,78,79} neural tube defects (defects of the brain, spine, or spinal cord);^{80,81} cardiovascular/heart defects; musculoskeletal defects, limb reduction defects; missing/extra digits; clubfoot; craniosynostosis (fused skull bones that may affect brain growth); facial defects; eye defects; gastrointestinal defects; gastroschisis (intestines and possibly other organs protrude from a hole beside the baby's navel); anal atresia; hernia; and undescended testes.⁷⁹
- Early developmental exposure to cigarette smoke can result in epigenetic changes in the lungs of the offspring which can be transferred to following generations, resulting in adult onset of respiratory disease.⁸² (Epigenetics is the study of biological mechanisms that switch genes on and off.) Some evidence shows an association between maternal smoking, early childhood exposure to secondhand smoke and the development of **emphysema in adulthood**. The findings suggest that the lungs may not recover completely from the effects of early-life exposure.^{83,84}
- Researchers have also found that adults exposed to tobacco smoke in utero had a more adverse **cardiovascular disease risk** profile.⁸⁵ There is even evidence that prenatal and postnatal secondhand smoke contributes to insulin resistance in children.⁸⁶
- One study found that maternal smoking during pregnancy is linked to high foetal testosterone (FT), which leads to an increased risk for **autism, ADHD, conduct disorder and antisocial behaviour**.⁸⁷ Numerous other studies have demonstrated a link between maternal smoking and ADHD,^{88,89,90} as well as new research linking secondhand smoke and ADHD.⁹¹
- A study examining causal relationships found that maternal smoking is associated with increased conduct problems and the externalising of problems in children. A limited association was found with hyperactivity and problems with peers.⁹²
- A population level study of children in Finland found that the risk of **psychiatric morbidity** was significantly higher in the children of mothers who smoked during pregnancy.⁹³
- Studies have found that smoking during pregnancy and exposure to secondhand smoke in early childhood were "quite strong" predictors of **conduct problems, anti-social behaviour** and crime later in life.^{94,95,96}
- This finding is supported by other studies exploring the relationship between maternal smoking during pregnancy and **behaviour problems** in childhood and adolescence,^{97,98,99,100,101,102} including **smoking** (particularly among girls)^{103,104} and other **substance use**.^{105,106,107}
- Maternal smoking has been associated with an increased risk of **learning difficulties**.¹⁰⁸
- Some evidence suggests that prenatal exposure to tobacco smoke may be associated with **benign breast disease** later in life.¹⁰⁹
- There is strong evidence that **childhood overweight and obesity** can be related to smoking during pregnancy.^{110,111,112,113,114} Two meta-analyses of 7 and 14 studies

respectively found that, for children of mothers who smoked during pregnancy, there was a 47-50% increase in the odds of being overweight in childhood.^{115,116}

Passive smoking and pregnancy

Secondhand tobacco smoke is a mixture of at least 4,000 chemical compounds,¹¹⁷ dozens of which are known or suspected reproductive toxins.¹³

Non-smoking women exposed to other people's tobacco smoke during pregnancy are more likely to have lower weight babies.¹¹⁸ On average, infants born to women exposed to secondhand smoke during pregnancy are 25-40g lighter than those born to women who are not exposed.¹³ Babies born to non-smoking women whose partners smoke have been found to weigh less than babies born to non-smoking couples.^{120,121} Exposure to second-hand smoke has also been significantly associated with preterm births.¹²² Other research suggests that non-smoking women who are exposed to second-hand smoke are at increased risk of difficulty becoming pregnant,^{13,123} of giving birth prematurely or stillbirth,^{124,125} of spontaneous abortion^{123,126} and of having a baby with congenital malformations.¹³

Additionally, some evidence suggests that female fertility can be damaged in utero if the woman's mother was exposed to secondhand smoke while pregnant.^{13,67,127}

It has been found that exposure to secondhand smoke can also be damaging in terms of successful pregnancy outcomes for women undergoing in vitro fertilization (IVF) or other Assisted Reproductive Technology (ART) treatment.^{128,129,130}

Attention deficit and hyperactivity have also been linked to secondhand smoke exposure in the home, in addition to maternal smoking during pregnancy.¹³¹

Exposure to parental secondhand smoke in the home and vehicle is strongly associated with middle ear disease in children.¹³²

Some studies show evidence of prenatal and postnatal secondhand smoke causing leukaemia,^{133,134} particularly acute lymphoblastic leukaemia.^{135,136,137}

Research on the reproductive effects of secondhand smoke exposure is relatively new compared to the effects of active smoking. More epidemiologic research is likely to reveal additional negative health effects, as well as the mechanisms whereby they occur and the dose-response relationships involved.¹³ Studies related to genetics and reproduction among smokers are also emerging and will continue to shed light on the processes involved.^{138,139}

Breast-feeding

Women who have quit smoking for at least a month are more apt to initiate breastfeeding.¹⁴⁰ In addition, women who quit smoking tend to continue breastfeeding for a longer period of time than those who continue to smoke.¹¹ A study of Canadian women found maternal smoking was negatively associated with 6 months exclusive breastfeeding.¹⁴² Women may not be aware that breastfeeding is still recommended by the National Health Service (NHS)¹⁴³ and the American Academy of Pediatrics,¹⁴⁴ among others, because of its beneficial effects on the baby, even if the mother continues to smoke.^{145,146} Health professionals should therefore combine smoking cessation and relapse prevention advice with lactation counselling to maximize success of smokers' efforts to initiate breastfeeding.^{147,148}

Smoking and oral contraception

Women who use combined oral contraceptives are at increased risk of heart disease. Because the risk of heart disease in young women is low, the benefits of using the pill generally outweigh

the risks for young women who do not smoke. Pill-users who smoke are also at risk of venous thromboembolism and arterial thrombosis (blood clots in the veins and arteries).^{149,150,151} It is therefore important that all women who take the contraceptive pill be advised not to smoke.

Smoking and the menopause

Smoking is associated with early onset of menopause^{152,153} with the natural menopause occurring up to two years earlier in smokers.¹⁵⁴ The likelihood of an earlier menopause is related to the number of cigarettes smoked, with those smoking more than ten cigarettes a day having an increased risk of an early menopause.¹⁵⁵

Stopping smoking may lower the risk of early menopause. While current smokers' risk of early menopause is twice that of non-smokers, in ex-smokers the risk is higher by just one-third. Research suggests that polycyclic aromatic hydrocarbons found in tobacco smoke can trigger premature egg cell death which may in turn lead to earlier menopause.¹⁵⁶ Another study suggests that chemicals in tobacco smoke alter endocrine function which in turn affects the release of pituitary hormones. This endocrine disruption is thought to contribute to adverse outcomes including earlier menopause.¹⁵⁷

Quitting smoking during pregnancy

More women quit smoking when they are pregnant than at any other time during their lives.¹⁵⁸ One study found that pregnant smokers were twice as likely to attempt to quit smoking as non-pregnant women¹⁵⁹ but surveys suggest that less than half of pregnant women smokers actually stop smoking during pregnancy. For example, in England, in 2015-16 17,743 pregnant women set a quit date with NHS Stop Smoking Services and 45% of them (7,914) successfully quit.¹⁶⁰ This is down from the 8,838 women who successfully quit in 2014-15.¹⁶¹

Because smoking poses a high risk of harm to both mother and foetus, it is important that pregnant women be supported to help them stop smoking at least for the duration of the pregnancy, but also postpartum. The NHS Pregnancy Smoking Helpline (0800 169 9 169) provides a call-back service that helps support women smokers throughout their pregnancy. However, support should come from the baby's father, family members and friends, as well as the health care system.

A 2009 Cochrane review found that interventions from health professionals reduced the proportion of women smoking in late pregnancy by about 6% overall.¹⁶²

The Department of Health has published guidance on stop smoking interventions in primary and secondary care which recommends establishing treatment pathways for all smokers including pregnant women.⁴⁰ The National Institute for Health and Clinical Excellence (NICE) has also published guidance on stopping smoking in pregnancy and following childbirth.¹⁶³

In March 2012, following a challenge by the Public Health Minister to investigate ways of further reducing the rate of smoking in pregnancy, the Smoking in Pregnancy Challenge Group was established to encourage health care professionals to work together towards this goal.¹⁶⁴ A follow-up report to measure progress was published in October 2015.¹⁶⁵

Smoking cessation interventions have been shown to reduce the number of newborns with low birth weight and preterm births.¹⁶² In one study, compared to ongoing heavy smoking, quitting was associated with a 299g increase in birth weight and going from heavy to light smoking was associated with a 199g increase in birth weight, while the babies of light smokers who quit entirely saw a 63g increase in birth weight.¹⁶⁶

As stopping smoking can be difficult, some women may need pharmacotherapy support. A review of the efficacy and safety of nicotine replacement therapy (NRT) in pregnancy found that NRT increased smoking cessation rates, measured in late pregnancy, by approximately 40%.¹⁶⁷ The authors found no evidence that NRT used for smoking cessation in pregnancy has either positive or negative impacts on birth outcomes.

A US review of studies found that NRT use significantly decreased the risk of preterm delivery and low birth weight compared to that of smokers and also found that NRT use does not appear to increase the risk of malformations.¹⁶⁸

Since NRT is metabolised up to 60% faster by pregnant women, higher doses of NRT may be needed.¹⁶⁷ Additional research is needed on which types and doses of NRT are the most effective in helping pregnant smokers quit, while also protecting the foetus.¹⁶⁷

Currently, the National Institute for Health and Care Excellence (NICE) recommends the use of NRT by pregnant women only if they have been unable to quit without NRT.¹⁶⁹

NICE does not recommend the use of either bupropion or varenicline by pregnant or breastfeeding women.¹⁶⁹

There is some research which suggests that financial incentives can increase quit rates among some pregnant smokers. One study found it to be the most effective intervention, particularly among low income women.¹⁷⁰

Even with pharmacotherapeutic, professional and social support, quitting is hard for some people due to their physical, psychological and social addictions to smoking. Many women will quit during their pregnancy, but the rate of relapse postpartum is high and this period should continue to be a focus for providing support to new mothers.¹⁷¹ One way that they can be supported is through initiatives to make homes smoke-free, thus protecting all family members from smoke exposure and reducing the chances of relapse.

Other resources

[Use of Electronic Cigarettes in Pregnancy: a guide for midwives and other healthcare professionals](#), Smoking in Pregnancy Challenge Group, 2016

References

- 1 Soares SR, Melo MA. [Cigarette smoking and reproductive function](#). Curr Opin Obstet Gynecol. 2008 Jun;20(3):281-91. doi: 10.1097/GCO.0b013e3282fc9c1e.
- 2 Dechanet C et al. [Effects of cigarette smoking on reproduction](#). Hum Reprod Update. 2011 Jan-Feb;17(1):76-95. doi: 10.1093/humupd/dmq033.
- 3 Dietz PM et al. [Infant morbidity and mortality attributable to prenatal smoking in the U.S.](#) Am J Prev Med. 2010 Jul;39(1):45-52. doi: 10.1016/j.amepre.2010.03.009.
- 4 Royal College of Physicians, Tobacco Advisory Group. Ch 3. [Effects of smoking on fetal and reproductive health](#). In: Passive smoking and children: A report by the Tobacco Advisory Group of the Royal College of Physicians. 2010 Mar.
- 5 Shiverick KT. Chapter 24 – Cigarette smoking and reproductive and developmental toxicity. In: Gupta RC, editor. Reproductive and Developmental Toxicology Burlington, MA: Elsevier; 2011. ISBN: 978-0-12-382032-7.
- 6 Soares SR et al. [Cigarette smoking affects uterine receptiveness](#). Human Reproduction. 2007 22(2):543-547; doi:10.1093/humrep/del394.
- 7 Einarson A, Riordan S. [Smoking in pregnancy and lactation: a review of risks and cessation strategies](#). Eur J Clin Pharmacol. 2009 Apr;65(4):325-30. doi: 10.1007/s00228-008-0609-0.
- 8 Anderson K, Nisenblat V, Norman R. [Lifestyle factors in people seeking infertility treatment - A review](#). Aust N Z J Obstet Gynaecol. 2010 Feb;50(1):8-20. doi: 10.1111/j.1479-828X.2009.01119.x.
- 9 Dechanet C et al. [Effects of cigarette smoking on embryo implantation and placentation and analysis of factors interfering with cigarette smoke effects \(Part II\)](#). [Article in French] Gynecol Obstet Fertil. 2011 Oct;39(10):567-74. doi: 10.1016/j.gyobfe.2011.07.023.
- 10 Lintsen A et al. [Effects of subfertility cause, smoking and body weight on the success rates of IVF](#). Hum Reprod. 2005 Jul;20(7):1867-75.
- 11 Neal M et al. [Sidestream smoking is equally as damaging as mainstream smoking on IVF outcomes](#). Hum Reprod. 2005 Sep;20(9):2531-5.
- 12 Waylen AL et al. [Effects of cigarette smoking upon clinical outcomes of assisted reproduction: a meta-analysis](#). Hum Reprod Update. 2009 Jan-Feb;15(1):31-44. doi: 10.1093/humupd/dmn046.
- 13 Meeker JD, Benedict MD. [Infertility, pregnancy loss and adverse birth outcomes in relation to maternal secondhand tobacco smoke exposure](#). Curr Womens Health Rev. 2013 Feb;9(1):41-49.
- 14 Cooper AR and Moley, KH. [Maternal tobacco use and its preimplantation effects on fertility: more reasons to stop smoking](#). Semin Reprod Med. 2008 Mar;26(2):204-12. doi: 10.1055/s-2008-1042959.
- 15 Ye X, Skjaerven R, Basso O et al. Hum Reprod. [In utero exposure to tobacco smoke and subsequent reduced fertility in females](#). 2010 Nov;25(11):2901-6. doi: 10.1093/humrep/deq235.
- 16 Ramlau-Hansen CH et al. [Is prenatal exposure to tobacco smoking a cause of poor semen quality? A follow-up study](#). Am J Epidemiol. 2007 Jun 15;165(12):1372-9.
- 17 Mamsen, LS et al. [Cigarette smoking during early pregnancy reduces the number of embryonic germ and somatic cells](#). Hum Reprod. 2010 Nov;25(11):2755-61. doi: 10.1093/humrep/deq215.
- 18 Hammadeh ME et al. [Protamine contents and P1/P2 ratio in human spermatozoa from smokers and non-smokers](#). Hum Reprod. 2010 Nov;25(11):2708-20. doi: 10.1093/humrep/deq226.
- 19 Sobinoff A et al. [Damaging legacy: maternal cigarette smoking has long-term consequences for male offspring fertility](#) Human Reproduction, 2014 Dec; 29(12):2719-35.
- 20 Virtanen HE et al. [Prenatal exposure to smoking and male reproductive health](#). Curr Opin Endocrinol Diabetes Obes. 2012 Jun;19(3):228-32. doi: 10.1097/MED.0b013e3283537cb8.
- 21 Jensen MS et al. [Lower sperm counts following prenatal tobacco exposure](#). Hum Reprod 2005; 20(9): 2559-66
- 22 Ramlau-Hansen CH et al. [Is smoking a risk factor for decreased semen quality? A cross-sectional analysis](#). Hum Reprod. 2007 Jan;22(1):188-96
- 23 Gaur DS, Talekar M, Pathak VP. [Effect of cigarette smoking on semen quality of infertile men](#). Singapore Med J. 2007 Feb;48(2):119-23.
- 24 Schilling K et al. [Prevalence of behaviour-related fertility disorders in a clinical sample: results of a pilot study](#). Arch Gynecol Obstet. 2012 Nov;286(5):1307-14. doi: 10.1007/s00404-012-2436-x.

- 25 Sofikitis N et al. [Effects of nicotine on sperm motility, membrane function and fertilizing capacity in vitro](#). Urol Res. 2000 Dec;28(6):370-5.
- 26 Ramlau-Hansen CH et al. [Is prenatal exposure to tobacco smoking a cause of poor semen quality? A follow-up study](#). Am J of Epidemiology 2007 165(12):1372-1379; doi:10.1093/aje/kwm032
- 27 Millett C et al. [Smoking and erectile dysfunction: findings from a representative sample of Australian men](#). Tob Control. 2006 Apr;15(2):136-9.
- 28 Meldrum DR et al. [Lifestyle and metabolic approaches to maximizing erectile and vascular health](#). Int J Impot Res. 2012 Mar-Apr;24(2):61-8. doi: 10.1038/ijir.2011.51.
- 29 Tostes RC et al. [Cigarette smoking and erectile dysfunction: focus on NO bioavailability and ROS generation](#). J Sex Med. 2008 Jun;5(6):1284-95. doi: 10.1111/j.1743-6109.2008.00804.x.
- 30 Grant P et al. [Erectile dysfunction in general medicine](#). Clin Med. 2013 Apr;13(2):136-40. doi: 10.7861/clinmedicine.13-2-136.
- 31 Heidelbaugh JJ. [Management of erectile dysfunction](#). Am Fam Physician. 2010 Feb 1;81(3):305-12
- 32 Flemming K et al. [Smoking in pregnancy: a systematic review of qualitative research of women who commence pregnancy as smokers](#). J Adv Nurs. 2013 May;69(5):1023-36. doi: 10.1111/jan.12066.
- 33 Department of Health. [Healthy lives, Healthy people: A Tobacco Control Plan for England](#). March 9, 2011.
- 34 [Smoking in Pregnancy Challenge Group](#)
- 35 Health and Social Care Information Centre. Statistical Release. [Statistics on women's smoking status at time of delivery: England. Quarter 4 2015/16](#). June 2016
- 36 Smoking in Pregnancy Challenge Group, [Smoking Cessation in Pregnancy: A review of the challenge](#), October 2015.
- 37 Vielwerth SE et al. [The impact of maternal smoking on fetal and infant growth](#). Early Hum Dev. 2007 Aug;83(8):491-5.
- 38 Agrawal A et al. [The effects of maternal smoking during pregnancy on offspring outcomes](#). Prev Med. 2010 Jan-Feb;50(1-2):13-8. doi: 10.1016/j.ypmed.2009.12.009.
- 39 McCowan L, Horgan RP. [Risk factors for small for gestational age infants](#). Best Pract Res Clin Obstet Gynaecol. 2009 Dec;23(6):779-93. doi: 10.1016/j.bpobgyn.2009.06.003.
- 40 Department of Health. [Local Stop Smoking Services: Service delivery and monitoring guidance 2011/12](#). March 14, 2011.
- 41 Juárez SP, Merlo J. [Revisiting the effect of maternal smoking during pregnancy on offspring birthweight: a quasi-experimental sibling analysis in Sweden](#). PLoS One. 2013 Apr 17;8(4):e61734. doi: 10.1371/journal.pone.0061734.
- 42 Samper MP et al. [Maternal active smoking and newborn body composition](#). Early Hum Dev. 2012 Mar;88(3):141-5. doi: 10.1016/j.earlhumdev.2011.07.015.
- 43 Varvarigou AA, Sotirios F, Nicholas BG. [Effect of prenatal tobacco smoke exposure on fetal growth potential](#). J Perinat Med. 2010 Nov;38(6):683-7. doi: 10.1515/JPM.2010.101.
- 44 Roza SJ et al. [Effects of maternal smoking in pregnancy on prenatal brain development](#). The Generation R Study. Eur J Neurosci. 2007 Feb;25(3):611-7.
- 45 Himes SK et al. [Prenatal tobacco exposure, biomarkers for tobacco in meconium, and neonatal growth outcomes](#). J Pediatr. 2013 May;162(5):970-5. doi: 10.1016/j.jpeds.2012.10.045.
- 46 Andersen MR et al. [Smoking cessation early in pregnancy and birth weight, length, head circumference, and endothelial nitric oxide synthase activity in umbilical and chorionic vessels: an observational study of healthy singleton pregnancies](#). Circulation. 2009 Feb 17;119(6):857-64. doi: 10.1161/CIRCULATIONAHA.107.755769.
- 47 British Medical Association, Board of Science; Education and Tobacco Control Resource Centre. [Smoking and reproductive life](#). The impact of smoking on sexual, reproductive and child health. London, BMA, 2004.
- 48 Gardosi J et al. [Stillbirth and infant mortality, West Midlands 1997-2005: Trends, Factors, Inequalities](#). West Midlands Perinatal Institute, 2007.
- 49 Abbott LC, Winzer-Serhan UH. [Smoking during pregnancy: lessons learned from epidemiological studies and experimental studies using animal models](#). Crit Rev Toxicol. 2012 Apr;42(4):279-303. doi: 10.3109/10408444.2012.658506.
- 50 Rogers JM. [Tobacco and pregnancy](#). Reprod Toxicol. 2009 Sep;28(2):152-60. doi: 10.1016/j.reprotox.2009.03.012.

- 51 Rubens D, Sarnat HB. [Sudden infant death syndrome: an update and new perspectives of etiology](#). *Handb Clin Neurol*. 2013;112:867-74. doi: 10.1016/B978-0-444-52910-7.00008-8.
- 52 Van Nguyen JM, Abenhaim HA. [Sudden Infant Death Syndrome](#): Review for the Obstetric Care Provider. *Am J Perinatol*. 2013 Jan 5.
- 53 McDonnell-Naughton M et al. [Maternal smoking and alcohol consumption during pregnancy as risk factors for sudden infant death](#). *Ir Med J*. 2012 Apr;105(4):105-8.
- 54 Liebrechts-Akkerman G et al. [Postnatal parental smoking: an important risk factor for SIDS](#). *Eur J Pediatr*. 2011 Oct;170(10):1281-91. doi: 10.1007/s00431-011-1433-6.
- 55 Lavezzi AM, Corna MF, Matturri L. [Ependymal alterations in sudden intrauterine unexplained death and sudden infant death syndrome: possible primary consequence of prenatal exposure to cigarette smoking](#). *Neural Dev*. 2010 Jul 19;5:17. doi: 10.1186/1749-8104-5-17
- 56 Behm I et al. [Increasing prevalence of smoke-free homes and decreasing rates of sudden infant death syndrome in the United States](#): an ecological association study. *Tob Control*. 2012 Jan;21(1):6-11. doi: 10.1136/tc.2010.041376.
- 57 Kyrklund-Blomberg NB, Granath F, Cnattingius S. [Maternal smoking and causes of very preterm birth](#). *Acta Obstet Gynecol Scand*. 2005 Jun;84(6):572-7.
- 58 Burke H et al. [Prenatal and passive smoke exposure and incidence of asthma and wheeze](#): systematic review and meta-analysis. *Pediatrics*. 2012 Apr;129(4):735-44. doi: 10.1542/peds.2011-2196.
- 59 Robison RG et al. [Maternal smoking during pregnancy, prematurity and recurrent wheezing in early childhood](#). *Pediatr Pulmonol*. 2012 Jul;47(7):666-73. doi: 10.1002/ppul.22501.
- 60 Duijts L et al. [Fetal exposure to maternal and paternal smoking and the risks of wheezing in preschool children](#): the Generation R Study. *Chest*. 2012 Apr;141(4):876-85. doi: 10.1378/chest.11-0112.
- 61 Gilliland F, Li Y, and Peters J. [Effects of Maternal Smoking during Pregnancy and Environmental Tobacco Smoke on Asthma and Wheezing in Children](#). *American Journal of Respiratory and Critical Care Medicine* Vol. 163, No. 2 (2001).
- 62 Noakes S et al. [Maternal smoking is associated with impaired neonatal Toll-like receptor \(TLR\) mediated immune responses](#). *Eur Respir J*. 2006 Oct;28(4):721-9.
- 63 Shenassa ED, Brown MJ. [Maternal smoking and infantile gastrointestinal dysregulation](#): the case of colic. *Pediatrics*. 2004 Oct;114(4):e497-505.
- 64 Canivet CA et al. [Infantile colic, maternal smoking and infant feeding at 5 weeks of age](#). *Scand J Public Health*. 2008 May;36(3):284-91. doi: 10.1177/1403494807086981.
- 65 Agrawal A et al. [The effects of maternal smoking during pregnancy on offspring outcomes](#). *Prev Med*. 2010 Jan-Feb;50(1-2):13-8. doi: 10.1016/j.ypmed.2009.12.009.
- 66 Boutwell BB, Beaver KM. [Maternal cigarette smoking during pregnancy and offspring externalizing behavioral problems](#): a propensity score matching analysis. *Int J Environ Res Public Health*. 2010 Jan;7(1):146-63. doi: 10.3390/ijerph7010146.
- 67 Meeker JD et al. [Maternal exposure to second-hand tobacco smoke and pregnancy outcome among couples undergoing assisted reproduction](#). *Human Reproduction* 2007; 22(2):337-345; doi:10.1093/humrep/del406.
- 68 Jaddoe VVW, de Ridder MAJ, Van Den Elzen APM. [Maternal smoking in pregnancy is associated with cholesterol development in the offspring](#): A 27-years follow-up study. *Atherosclerosis*. 2008 Jan;196(1):42-8.
- 69 Ayer JG et al. [Maternal cigarette smoking is associated with reduced high-density lipoprotein cholesterol in healthy 8-year-old children](#). *Eur Heart J*. 2011 Oct;32(19):2446-53. doi: 10.1093/eurheartj/ehr174.
- 70 Wen X, Triche EW, Hogan JW. [Birth Weight and Adult Hypercholesterolemia](#): Subgroups of small-for-gestational-age based on maternal smoking status during pregnancy. *Epidemiology*. 2010 Nov;21(6):786-90. doi: 10.1097/EDE.0b013e3181f20990.
- 71 Stjernfeld M et al. [Maternal smoking during pregnancy and risk of childhood cancer](#). *The Lancet* Volume 327, Issue 8494, 14 June 1986.
- 72 Bhattacharya S. et al. [Maternal and perinatal risk factors for childhood cancer: record linkage study](#). *BMJ Open* 2014; 4:e003656 doi:10.1136.
- 73 Heck J. et al. [Smoking in pregnancy and risk of cancer among young children: A population-based study](#). *International Journal of Cancer* 2016; Volume 139, Issue 3.
- 74 Ellingson JM et al. [Disentangling the relationships between maternal smoking during pregnancy and co-occurring risk factors](#). *Psychol Med*. 2012 Jul;42(7):1547-57. doi: 10.1017/S0033291711002534

- 75 Leonardi-Bee J, Britton J, Venn A. [Secondhand smoke and adverse fetal outcomes in nonsmoking pregnant women](#): a meta-analysis. *Pediatrics*. 2011 Apr;127(4):734-41. doi: 10.1542/peds.2010-3041.
- 76 Lebbby KD, Tan F, Brown CP. [Maternal factors and disparities associated with oral clefts](#). *Ethn Dis*. 2010 Winter;20(1 Suppl 1):S1-146-9.
- 77 Li L et al. [Biological and epidemiological evidence of interaction of infant genotypes at Rs7205289 and maternal passive smoking in cleft palate](#). *Am J Med Genet A*. 2011 Dec;155A(12):2940-8. doi: 10.1002/ajmg.a.34254.
- 78 Shi M, Wehby GL, Murray JC. [Review on genetic variants and maternal smoking in the etiology of oral clefts and other birth defects](#). *Birth Defects Res C Embryo Today*. 2008 Mar;84(1):16-29. doi: 10.1002/bdrc.20117.
- 79 Hackshaw A, Rodeck C, Boniface S. [Maternal smoking in pregnancy and birth defects](#): a systematic review based on 173 687 malformed cases and 11.7 million controls. *Hum Reprod Update*. 2011 Sep-Oct;17(5):589-604. doi: 10.1093/humupd/dmr022.
- 80 Suarez L et al. [Maternal exposures to cigarette smoke, alcohol, and street drugs and neural tube defect occurrence in offspring](#). *Matern Child Health J*. 2008 May;12(3):394-401.
- 81 Suarez L et al. [Maternal smoking, passive tobacco smoke, and neural tube defects](#). *Birth Defects Res A Clin Mol Teratol*. 2011 Jan;91(1):29-33. doi: 10.1002/bdra.20743.
- 82 Harding R, Maritz G. [Maternal and fetal origins of lung disease in adulthood](#). *Semin Fetal Neonatal Med*. 2012 Apr;17(2):67-72. doi: 10.1016/j.siny.2012.01.005.
- 83 Lovasi GS et al. [Association of environmental tobacco smoke exposure in childhood with early emphysema in adulthood among nonsmokers](#). *Am J Epidemiol*. 2010 Jan 1;171(1):54-62. doi: 10.1093/aje/kwp358.
- 84 Beyer D, Mitfessel H, Gillissen A. [Maternal smoking promotes chronic obstructive lung disease in the offspring as adults](#). *Eur J Med Res*. 2009 Dec 7;14 Suppl 4:27-31.
- 85 Power C, Atherton K, Thomas C. [Maternal smoking in pregnancy, adult adiposity and other risk factors for cardiovascular disease](#). *Atherosclerosis*. 2010 Aug;211(2):643-8. doi: 10.1016/j.atherosclerosis.2010.03.015.
- 86 Thiering E et al. [Prenatal and postnatal tobacco smoke exposure and development of insulin resistance in 10 year old children](#). *Int J Hyg Environ Health*. 2011 Sep;214(5):361-8. doi: 10.1016/j.ijheh.2011.04.004.
- 87 Rizwan S, Manning JT, Brabin BJ. [Maternal smoking during pregnancy and possible effects of in utero testosterone](#): Evidence from the 2D: 4D finger length ratio. *Early Hum Dev*. 2007 Feb;83(2):87-90.
- 88 Motlagh MG et al. [Adverse effects of heavy prenatal maternal smoking on attentional control in children with ADHD](#). *J Atten Disord*. 2011 Oct;15(7):593-603. doi: 10.1177/1087054710374576.
- 89 Nomura Y, Marks DJ, Halperin JM. [Prenatal exposure to maternal and paternal smoking on attention deficit hyperactivity disorders symptoms and diagnosis in offspring](#). *J Nerv Ment Dis*. 2010 Sep;198(9):672-8. doi: 10.1097/NMD.0b013e3181ef3489.
- 90 Obel C et al. [Is maternal smoking during pregnancy a risk factor for hyperkinetic disorder?--Findings from a sibling design](#). *Int J Epidemiol*. 2011 Apr;40(2):338-45. doi: 10.1093/ije/dyq185.
- 91 Max W, Sung HY, Shi Y. [Attention deficit hyperactivity disorder among children exposed to secondhand smoke](#): a logistic regression analysis of secondary data. *Int J Nurs Stud*. 2013 Jun;50(6):797-806. doi: 10.1016/j.ijnurstu.2012.10.002.
- 92 Brion M. [Maternal Smoking and Child Psychological Problems: Disentangling Causal and Noncausal Effects](#). *Paediatrics* 2010; 126 (1).
- 93 Ekblad M et al. [Prenatal smoking exposure and the risk of psychiatric morbidity into young adulthood](#). *Arch Gen Psychiatry*. 2010 Aug;67(8):841-9. doi: 10.1001/archgenpsychiatry.2010.92.
- 94 Murray J et al. [Very early predictors of conduct problems and crime](#): results from a national cohort study. *J Child Psychol Psychiatry*. 2010 Nov;51(11):1198-207. doi: 10.1111/j.1469-7610.2010.02287.x.
- 95 Paradis AD et al. [Maternal smoking during pregnancy and criminal offending among adult offspring](#). *J Epidemiol Community Health*. 2011 Dec;65(12):1145-50. doi: 10.1136/jech.2009.095802.
- 96 Cornelius MD, Goldschmidt L, Day NL. [Prenatal cigarette smoking: Long-term effects on young adult behavior problems and smoking behavior](#). *Neurotoxicol Teratol*. 2012 Nov-Dec;34(6):554-9. doi: 10.1016/j.ntt.2012.09.003.
- 97 Bryon MJ et al. [Maternal smoking and child psychological problems](#): disentangling causal and noncausal effects. *Pediatrics*. 2010 Jul;126(1):e57-65. doi: 10.1542/peds.2009-2754.
- 98 Liu J et al. [Mother's environmental tobacco smoke exposure during pregnancy and externalizing behavior problems in children](#). *Neurotoxicology*. 2013 Jan;34:167-74. doi: 10.1016/j.neuro.2012.11.005.

- 99 Cornelius MD et al. [Effects of prenatal cigarette smoke exposure on neurobehavioral outcomes in 10-year-old children of adolescent mothers](#). *Neurotoxicol Teratol.* 2011 Jan-Feb;33(1):137-44. doi: 10.1016/j.ntt.2010.08.006.
- 100 Cornelius MD et al. [Long-term effects of prenatal cigarette smoke exposure on behavior dysregulation among 14-year-old offspring of teenage mothers](#). *Matern Child Health J.* 2012 Apr;16(3):694-705. doi: 10.1007/s10995-011-0766-0.
- 101 Piper BJ, Gray HM, Birkett MA. [Maternal smoking cessation and reduced academic and behavioral problems in offspring](#). *Drug Alcohol Depend.* 2012 Feb 1;121(1-2):62-7. doi: 10.1016/j.drugalcdep.2011.08.004.
- 102 Huijbregts SC et al. [Neurobiological and behavioral stress reactivity in children prenatally exposed to tobacco](#). *Psychoneuroendocrinology.* 2011 Jul;36(6):913-8. doi: 10.1016/j.psyneuen.2010.12.008.
- 103 Rydell M et al. [Prenatal exposure to tobacco and future nicotine dependence: population-based cohort study](#). *Br J Psychiatry.* 2012 Mar;200(3):202-9. doi: 10.1192/bjp.bp.111.100123.
- 104 O'Callaghan FV, Al Mamun A, O'Callaghan M et al. [Maternal smoking during pregnancy predicts nicotine disorder \(dependence or withdrawal\) in young adults - a birth cohort study](#). *Aust N Z J Public Health.* 2009; 33(4):371-7
- 105 Monshouwer K et al. [Prenatal smoking exposure and the risk of behavioral problems and substance use in adolescence: the TRAILS study](#). *Eur Addict Res.* 2011;17(6):342-50. doi: 10.1159/000334507.
- 106 Goldschmidt L, Cornelius MD, Day NL. [Prenatal cigarette smoke exposure and early initiation of multiple substance use](#). *Nicotine Tob Res.* 2012 Jun;14(6):694-702. doi: 10.1093/ntr/ntr280.
- 107 Nomura Y, Gilman SE, Buka SL. [Maternal smoking during pregnancy and risk of alcohol use disorders among adult offspring](#). *J Stud Alcohol Drugs.* 2011 Mar;72(2):199-209.
- 108 Anderko L, Braun J, Auinger P. [Contribution of tobacco smoke exposure to learning disabilities](#). *J Obstet Gynecol Neonatal Nurs.* 2010 Jan-Feb;39(1):111-7. doi: 10.1111/j.1552-6909.2009.01093.x.
- 109 Liu T et al. [Prenatal exposure to cigarette smoke and benign breast disease](#). *Epidemiology.* 2010 Sep;21(5):736-43. doi: 10.1097/EDE.0b013e3181e9c118.
- 110 Braun JM et al. [Prenatal environmental tobacco smoke exposure and early childhood body mass index](#). *Paediatr Perinat Epidemiol.* 2010 Nov;24(6):524-34. doi: 10.1111/j.1365-3016.2010.01146.x.
- 111 Mangrio E, Lindström M, Rosvall M. [Early life factors and being overweight at 4 years of age among children in Malmö, Sweden](#). *BMC Public Health.* 2010 Dec 15;10:764. doi: 10.1186/1471-2458-10-764.
- 112 Chen H et al. [Cigarette smoking and brain regulation of energy homeostasis](#). *Front Pharmacol.* 2012;3:147. doi: 10.3389/fphar.2012.00147.
- 113 Raum E et al. [Tobacco smoke exposure before, during, and after pregnancy and risk of overweight at age 6](#). *Obesity (Silver Spring).* 2011 Dec;19(12):2411-7. doi: 10.1038/oby.2011.129.
- 114 Toshihiro I. et al. [Maternal smoking during pregnancy and offspring obesity: Meta-analysis](#). *Pediatrics International,* 52: 94–99. doi:10.1111/j.1442-200X.2009.02883.
- 115 Weng SF et al. [Systematic review and meta-analyses of risk factors for childhood overweight identifiable during infancy](#). *Arch Dis Child.* 2012 Dec;97(12):1019-26. doi: 10.1136/archdischild-2012-302263.
- 116 Oken E et al. [Maternal smoking during pregnancy and child overweight: systematic review and meta-analysis](#). *Int J Obes (Lond).* 2008 Feb;32(2):201-10. doi: 10.1038/sj.ijo.0803760.
- 117 [The health consequences of involuntary exposure to tobacco smoke](#): a report of the Surgeon General. U.S. Dept. of Health and Human Services, Centers for Disease Control and Prevention, Coordinating Center for Health Promotion, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, [2006]
- 118 Crane JM et al. *BJOG.* [Effects of environmental tobacco smoke on perinatal outcomes: a retrospective cohort study](#). 2011 Jun;118(7):865-71. doi: 10.1111/j.1471-0528.2011.02941.x.
- 119 Lee, J. et al. [Influence of maternal environmental tobacco smoke exposure assessed by hair nicotine levels on birth weight](#). *Asian Pac J Cancer Prev.* 2015;16(7):3029-34.
- 120 Khader YS et al. [The association between second hand smoke and low birth weight and preterm delivery](#). *Matern Child Health J.* 2011 May;15(4):453-9. doi: 10.1007/s10995-010-0599-2.
- 121 Salmasi G et al. [Environmental tobacco smoke exposure and perinatal outcomes: a systematic review and meta-analyses](#). *Acta Obstet Gynecol Scand.* 2010;89(4):423-41. doi: 10.3109/00016340903505748.
- 122 Mojibyan M. et al [Exposure to second-hand smoke during pregnancy and preterm delivery](#). *International Journal of High Risk Behaviours and Addiction* 2013 Mar 12. doi: 10.5812/ijhrba.7630.

- 123 Peppone LJ et al. [Associations between adult and childhood secondhand smoke exposures and fecundity and fetal loss among women who visited a cancer hospital](#). *Tob Control* 2009; 18(2): 115-20.
- 124 Subramoney S, d'Espaignet ET, Gupta PC. [Higher risk of stillbirth among lower and middle income women who do not use tobacco, but live with smokers](#). *Acta Obstet Gynecol Scand*. 2010;89(4):572-7. doi: 10.3109/00016341003801656.
- 125 Nieuwenhuijsen MJ et al. [Environmental risk factors of pregnancy outcomes: a summary of recent meta-analyses of epidemiological studies](#). *Environ Health*. 2013 Jan 15;12:6. doi: 10.1186/1476-069X-12-6.
- 126 George L et al. [Environmental tobacco smoke and risk of spontaneous abortion](#). *Epidemiology*. 2006 Sep;17(5):500-5.
- 127 Ye X et al. [In utero exposure to tobacco smoke and subsequent reduced fertility in females](#). *Hum Reprod* 2010; 25(11): 2901-6.
- 128 Penzias AS. [Recurrent IVF failure: other factors](#). *Fertil Steril*. 2012 May;97(5):1033-8. doi: 10.1016/j.fertnstert.2012.03.017.
- 129 Benedict MD et al. [Secondhand tobacco smoke exposure is associated with increased risk of failed implantation and reduced IVF success](#). *Hum Reprod*. 2011 Sep;26(9):2525-31. doi: 10.1093/humrep/der226.
- 130 Anderson K, Nisenblatt V, Norman R. [Lifestyle factors in people seeking infertility treatment](#) - A review. *Aust N Z J Obstet Gynaecol*. 2010 Feb;50(1):8-20. doi: 10.1111/j.1479-828X.2009.01119.x.
- 131 Tiesler CM et al. [Passive smoking and behavioural problems in children](#): results from the LISApplus prospective birth cohort study. *Environ Res*. 2011 Nov;111(8):1173-9. doi: 10.1016/j.envres.2011.06.011.
- 132 Jones LL et al. [Parental smoking and the risk of middle ear disease in children](#): a systematic review and meta-analysis. *Arch Pediatr Adolesc Med*. 2012 Jan;166(1):18-27. doi: 10.1001/archpediatrics.2011.158.
- 133 Chang JS. [Parental smoking and childhood leukemia](#). *Methods Mol Biol*. 2009;472:103-37. doi: 10.1007/978-1-60327-492-0_5.
- 134 (US) Department of Health and Human Services. [Children and Secondhand Smoke Exposure](#). Excerpts from The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General. Excerpts from Chapter 5 - Reproductive and Developmental Effects from Exposure to Secondhand Smoke. DHHS, Centers for Disease Control and Prevention, Office on Smoking and Health, 2007.
- 135 Milne E et al. [Parental prenatal smoking and risk of childhood acute lymphoblastic leukemia](#). *Am J Epidemiol*. 2012 Jan 1;175(1):43-53. doi: 10.1093/aje/kwr275.
- 136 Liu R, Zhang L, McHale CM, Hammond SK. [Paternal smoking and risk of childhood acute lymphoblastic leukemia](#): systematic review and meta-analysis. *J Oncol*. 2011;2011:854584. doi: 10.1155/2011/854584.
- 137 Metayer C et al. [Tobacco Smoke Exposure and the Risk of Childhood Acute Lymphoblastic and Myeloid Leukemias by Cytogenetic Subtype](#). *Cancer Epidemiol Biomarkers Prev*. 2013 Jul 12. [Epub ahead of print]
- 138 Knopik VS et al. [The epigenetics of maternal cigarette smoking during pregnancy and effects on child development](#). *Dev Psychopathol*. 2012 Nov;24(4):1377-90. doi: 10.1017/S0954579412000776.
- 139 Suter M et al. [Genetic and epigenetic influences associated with intrauterine growth restriction due to in utero tobacco exposure](#). *Pediatr Endocrinol Rev*. 2010 Dec;8(2):94-102.
- 140 Collins BN, DiSantis KI, Nair US. [Longer previous smoking abstinence relates to successful breastfeeding initiation among underserved smokers](#). *Breastfeed Med*. 2011 Dec;6(6):385-91. doi: 10.1089/bfm.2010.0076.
- 141 Higgins TM et al. [Effects of cigarette smoking cessation on breastfeeding duration](#). *Nicotine Tob Res*. 2010 May;12(5):483-8. doi: 10.1093/ntr/ntq031.
- 142 Al-Sahab B. et al. [Prevalence and predictors of 6-month exclusive breastfeeding among Canadian women: a national survey](#). *BMC Pediatrics*201010:20 DOI: 10.1186/1471-2431-10-20
- 143 National Health Service. NHS choices: Your health your choices. [Healthy lifestyle and breastfeeding](#). [Accessed 22 Aug 2016]
- 144 American Academy of Pediatrics. [Policy statement: Breastfeeding and the use of human milk](#). *Pediatrics*. 2005; 15(2):496–506.
- 145 Bailey BA, Wright HN. [Breastfeeding initiation in a rural sample](#): predictive factors and the role of smoking. *J Hum Lact*. 2011 Feb;27(1):33-40. doi: 10.1177/0890334410386955.
- 146 Dorea JG. [Maternal smoking and infant feeding](#): breastfeeding is better and safer. *Matern Child Health J*. 2007 May;11(3):287-91.
- 147 Collins BN, DiSantis KI, Nair US. [Longer previous smoking abstinence relates to successful breastfeeding initiation among underserved smokers](#). *Breastfeed Med*. 2011 Dec;6(6):385-91. doi: 10.1089/bfm.2010.0076.

- 148 Weiser TM et al. [Association of maternal smoking status with breastfeeding practices](#): Missouri, 2005. Pediatrics. 2009 Dec;124(6):1603-10. doi: 10.1542/peds.2008-2711.
- 149 Lalude OO. [Risk of cardiovascular events with hormonal contraception](#): insights from the Danish cohort study. Curr Cardiol Rep. 2013 Jul;15(7):374. doi: 10.1007/s11886-013-0374-2.
- 150 Pomp ER, Rosendaal FR, Doggen CJ. [Smoking increases the risk of venous thrombosis and acts synergistically with oral contraceptive use](#). Am J Hematol. 2008 Feb;83(2):97-102.
- 151 Shufelt CL, Bairey Merz CN. [Contraceptive hormone use and cardiovascular disease](#). J Am Coll Cardiol. 2009;53:221–31.
- 152 Saraç F, Oztekin K, Celebi G. [Early menopause association with employment, smoking, divorced marital status and low leptin levels](#). Gynecol Endocrinol. 2011 Apr;27(4):273-8. doi: 10.3109/09513590.2010.491165.
- 153 Sun L et al. [Meta-analysis suggests that smoking is associated with an increased risk of early natural menopause](#). Menopause. 2012 Feb;19(2):126-32. doi: 10.1097/gme.0b013e318224f9ac.
- 154 Office on Smoking and Health (US). [Women and Smoking: A Report of the Surgeon General](#). Atlanta (GA): Centers for Disease Control and Prevention (US); 2001 Mar.
- 155 Kinney A, Klinec J, Levin B. [Alcohol, caffeine and smoking in relation to age at menopause](#). Maturitas. Volume 54, Issue 1, 20 April 2006, Pages 27–38.
- 156 Matikainen T et al. [Aromatic hydrocarbon receptor-driven Bax gene expression is required for premature ovarian failure caused by biohazardous environmental chemicals](#). Nat Genet. 2001 Aug;28(4):355-60.
- 157 Windham G et al. [Cigarette smoking and effects on hormone function in premenopausal women](#). Environ Health Perspect. 2005 Oct;113(10):1285-90.
- 158 Murin S, Ruffin R, Bilello K. [Smoking and smoking cessation in pregnancy](#). Clin Chest Med. 2011 Mar;32(1):75-91, viii. doi: 10.1016/j.ccm.2010.11.004.
- 159 Chen X et al. [Age of smoking onset as a predictor of smoking cessation during pregnancy](#). Am J Health Behav. 2006;30: 247–258.
- 160 [Statistics on NHS Stop Smoking Services: England, April 2015 – March 2016](#). NHS Digital 18 August 2016.
- 161 [Statistics on NHS Stop Smoking Services: England, April 2014 – March 2015](#). Health and Social Care Information Centre, 19 August 2015.
- 162 Lumley J et al. [Interventions for promoting smoking cessation during pregnancy](#). Cochrane Database Syst Rev. 2009 Jul 8;(3):CD001055. doi: 10.1002/14651858.CD001055.pub3.
- 163 National Institute for Health and Care Excellence. [Quitting smoking in pregnancy and following childbirth](#): NICE public health guidance 26. 2010 Jun.
- 164 Action on Smoking and Health (ASH) [Smoking cessation in pregnancy: A call to action](#). Lowry C and Scammell K (editors.). June 2013
- 165 Action on Smoking and Health (ASH) [Smoking cessation in pregnancy: A review of the challenge](#). October 2015.
- 166 Benjamin-Garner R, Stotts A. [Impact of smoking exposure change on infant birth weight among a cohort of women in a prenatal smoking cessation study](#). Nicotine Tob Res. 2013 Mar;15(3):685-92. doi: 10.1093/ntr/nts184.
- 167 Coleman T et al. [Pharmacological interventions for promoting smoking cessation during pregnancy](#). Cochrane Database Syst Rev. 2012 Sep 12;9:CD010078. doi: 10.1002/14651858.CD010078.
- 168 Forinash AB et al. [Nicotine replacement therapy effect on pregnancy outcomes](#). Ann Pharmacother. 2010 Nov;44(11):1817-21. doi: 10.1345/aph.1P279.
- 169 National Institute for Health and Clinical Evidence. [Quitting smoking in pregnancy and following childbirth](#): NICE public health guidance 26. 2010 Jun.
- 170 Higgins St et al. [Financial incentives for smoking cessation among pregnant and newly postpartum women](#). Prev Med. 2012 Nov;55 Suppl:S33-40. doi: 10.1016/j.ypmed.2011.12.016.
- 171 Von Kohorn I et al. [A qualitative study of postpartum mothers' intention to smoke](#). Birth. 2012;39(1):65-9. doi: 10.1111/j.1523-536X.2011.00514.x.

