


Core20: Above-average smoking rates among the most deprived reduces their healthy life expectancy and increases pressure on the NHS



In England a third of all smokers live in the most deprived two deciles.¹ In NHS Lincolnshire ICB there are 83,100 smokers and 22% of people in routine and manual occupations smoke. Smoking costs your ICB £25.7m a year.²

Annually in your ICB smoking causes:

- 7,252 hospital admissions.³
- 1,122 premature deaths.⁴

Additional impact on communities in your ICB:

- 41,309 smoking households live in poverty.⁵
- 3,801 people out of work due to smoking.⁶
- 12,032 people receive informal care from friends and family because of smoking.⁷

PLUS: The most deprived groups have the highest smoking rates






National smoking rates among:

- People who are homeless (77%).⁸
- People entering prison (80%).⁹
- 11–16-year-olds with a mental disorder (22%).¹⁰
- People in social housing (26%).¹¹

Smoking rates for those receiving addiction treatment in your ICB:

- Those receiving treatment for opioid addiction (51%).¹²
- Those receiving treatment for alcohol addiction (34%).¹³

5: Five clinical areas of focus are all impacted by smoking

| 1. Maternity | 2. Severe Mental Illness | 3. Chronic respiratory illness | 4. Early cancer diagnosis | 5. Hypertension |
|--|---|---|---|---|
|  Smoking is the leading modifiable risk factor for poor birth outcomes. In your ICB 15% ¹⁴ of women smoke at time of delivery, 931 women annually. ¹⁵ |  Smoking is the leading cause of the 10-20 year reduction in life expectancy for people with serious mental illness (SMI). In your ICB 40% of people with SMI smoke. ¹⁶ |  Around 86% of all chronic obstructive pulmonary disease (COPD) deaths are caused by smoking. In your ICB 348 people a year die from COPD. ¹⁷ |  Smoking is the leading preventable cause of cancer responsible for 27% of cancer deaths. In your ICB 497 people a year die from cancer caused by smoking. ¹⁸ |  Smoking cessation is embedded in NICE guidelines on hypertension because smokers' CVD risk is double that of non-smokers. Nationally 9,300 people a year die from CVD caused by smoking. ¹⁹ |
| Find out more | Find out more | Find out more | Find out more | Find out more |

References

- [1] ONS. [Deprivation and the impact on smoking prevalence](#). 2023
- [2] [ASH Ready ICB Reckoner](#) 2023
- [3] [Tobacco dependence treatment services: delivery model \(2021\)](#) Admissions data from Hospital Episode Statistics (HES); Office for National Statistics (ONS) - mid-year population estimates; Smoking prevalence data from Annual Population Survey; and relative risks from the Royal College of Physician's Report 'Hiding in Plain Sight'. Data - OHID
- [4] [Smoking attributable mortality \(new method\)](#). 2017 – 19 Directly standardised rate - per 100,000 Local Tobacco Control Profiles - Data - OHID
- [5] H Reed (2021), [Estimates of poverty in the UK adjusted for expenditure on tobacco](#) – 2021 update.
- [6] [H Reed \(2020\), The impact of smoking history on employment prospects, earnings and productivity](#): an analysis using UK panel data.
- [7] H Reed (2021), [The costs of smoking to the social care system and related costs for older people in England](#): 2021 revision.
- [8] Homeless Link. [The Unhealthy State of Homelessness](#). Health audit results 2014.
- [9] O'Moore E. [Successfully delivering smokefree prisons across England and Wales](#). July 2018.
- [10] Marcheselli F, Brodie E, Si N, Pearce N, McManus S, Sadler K, et al. [Mental Health of Children and Young People in England, 2017](#) . NHS Digital; 2018.
- [11] [Local Tobacco Profiles](#) – Data – OHID: Housing tenure
- [12] [Smoking prevalence in adults \(18+\) admitted to treatment for substance misuse \(NDTMS\)](#) - all opiates. 2019/20 Local Tobacco Control Profiles - Data - OHID
- [13] [Smoking prevalence in adults \(18+\) admitted to treatment for substance misuse \(NDTMS\)](#) - alcohol. 2019/20 Local Tobacco Control Profiles - Data - OHID
- [14] [Smoking status at time of delivery](#). (2021) Calculated by PHE from the NHS Digital return on Smoking Status At Time of delivery (SATOD) Local Tobacco Control Profiles - Data - OHID
- [15] [Smoking status at time of delivery](#). (2021) Calculated by PHE from the NHS Digital return on Smoking Status At Time of delivery (SATOD) Local Tobacco Control Profiles - Data - OHID
- [16] [Smoking prevalence in adults \(18+\) with serious mental illness \(SMI\)](#) (2016) Local Tobacco Control Profiles - Data - OHID
- [17] [Mortality rate from chronic obstructive pulmonary disease](#) (3 year range) 2017 – 19 Directly standardised rate - per 100,000 Local Tobacco Control Profiles - Data - OHID
- [18] [Smoking attributable deaths from Cancer](#) (new method). 2017 – 19 Directly standardised rate - per 100,000 Local Tobacco Control Profiles - Data - OHID
- [19] [Smoking attributable deaths from heart disease](#) (new method). 2017 – 19 Directly standardised rate - per 100,000 Local Tobacco Control Profiles - Data - OH